Subject: Physics





Name of the Student:

Max. Marks: 22 Marks Time: 22 Minutes

Mark Schemes

## Q1.

(a) diagram to show: both focal points coinciding and labelled, with  $f_o > f_e$  (1) centre ray straight through objective, rays crossing at focal plane and proceeding to eyepiece (1) rays refracted at eyepiece and emerge parallel to construction line (1)

(b) (i)  $(f_o + f_e = 3.5, \text{ and } f_o / f_e = 100)$  estimate  $f_o \approx 3.5$  m and  $f_e \approx 0.035$  m (1)

(ii) (use of 
$$M = \frac{\alpha}{\alpha}$$
 gives)  $\alpha = \frac{4.0 \times 10^{-3}}{100} = 4.0 \times 10^{-5}$  (rad) (1) (use of  $\alpha = \frac{D}{r}$  gives)  $D = 4.0 \times 10^{-5} \times 1.3 \times 10^{9} = 5.2 \times 10^{4}$  km (1) (allow C.E. for value of  $\alpha$ )

(c) no chromatic aberration - mirrors do not refract light (1) no spherical aberration - use of parabolic mirror (1) no distortion - mirror can be supported more strongly (1) better resolving power or greater brightness - mirrors can be larger (1) more light gets through (image brighter) - lens absorbs more light (1) (any two)

[8]

3

3

2

**Q2**.

(a) (i) (use of 
$$\lambda_{\text{max}}T = 0.0029$$
 gives)  $\lambda = \frac{0.0029}{6000}$  (1) 
$$= 4.8 \times 10^{-7} \text{ m (1)}$$

- (ii) values on axis: 0.5 1.0 1.5 2.0 (1)
- (iii) similar shaped curve with peak shifted to right (1)

max 4

(b) (i) <u>difference</u> in absolute magnitude = 5 **(1)** corresponds to × 100 difference in brightness, some reference to absolute scale **(1)** 

Arcturus lower absolute magnitude, therefore brighter (1)

(= 200)

(ii) (use of 
$$P = \sigma A T^4$$
 gives) 
$$\frac{P_A}{P_S} = 100 = \frac{A_A T_A^4}{A_A T_S^4}$$
(1) 
$$\frac{A_A}{A_S} = 100 \times \left(\frac{6000}{5000}\right)^4$$
(1)

max 4

[8]

Q3.

(a) (i) 
$$\Delta \lambda = \frac{\lambda v}{c}$$
 (1)

(ii) 
$$\Delta \lambda = -\frac{\lambda v}{c}$$
 (1)

**(2)** 

(b) (i) total difference in wavelength = 
$$\frac{2\lambda v}{c}$$
 (1)

$$v = \frac{7.8 \times 10^{-12} \times 3.0 \times 10^{8}}{589 \times 10^{-9} \times 2} = 1986 \text{ [or } 2.0 \times 10^{3} \text{] m s}^{-1} \text{ (1)}$$

(ii) 
$$\omega = \frac{v}{r} = \frac{1986}{7.0 \times 10^8}$$
 (1)

= 
$$2.8 \times 10^{-6} \text{ rad s}^{-1}$$
 (1)

**(4)** [6]