

Name of the Student: \_\_\_\_\_

Max. Marks : 20 Marks

Time : 20 Minutes

**Q1.**

Atoms are different sizes.

One of the heaviest naturally occurring stable elements is lead.

Two of its isotopes are lead-206 ( $^{206}_{82}\text{Pb}$ ) and lead-208 ( $^{208}_{82}\text{Pb}$ ).

(a) (i) What is meant by 'isotopes'?

---

---

---

---

(2)

(ii) How many protons are in the nucleus of a  $^{206}_{82}\text{Pb}$  atom?

---

(1)

(iii) How many neutrons are in the nucleus of a  $^{206}_{82}\text{Pb}$  atom?

---

(1)

(b) A nucleus can be accelerated in a particle accelerator and directed at a large nucleus. This produces a heavy nucleus that will decay after a short time.


This is shown in **Figure 1**.

**Figure 1**



- (i) In 1984, nuclei of iron (Fe) were directed at nuclei of lead (Pb). This produced nuclei of hassium (Hs).

Complete the equation for this reaction by writing numbers in the empty boxes.



(3)

- (ii) Use the correct answer from the box to complete the sentence.

an electron	a proton	a neutron
-------------	----------	-----------

The particle **X** in part (b)(i) is \_\_\_\_\_.

(1)

- (iii) After acceleration the iron nuclei travel at a steady speed of one-tenth of the speed of light.

The speed of light is  $3.00 \times 10^8$  m/s.

Calculate the time taken for the iron nuclei to travel a distance of 12 000 m.

\_\_\_\_\_

\_\_\_\_\_

Time taken = \_\_\_\_\_ s

(2)

- (iv) Linear accelerators, in which particles are accelerated in a straight line, are **not** used for these experiments. Circular particle accelerators are used.

Suggest why.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

(3)

(c) Hassium-265 ( $^{265}_{108}\text{Hs}$ ) decays by alpha emission with a half-life of 0.002 seconds.

(i) What is meant by 'half-life'?

Tick (✓) **two** boxes.

	Tick (✓)
The average time for the number of nuclei to halve	
The time for count rate to be equal to background count	
The time for background count to halve	
The time for count rate to halve	

(2)

(ii) Complete the equation for the decay of Hs-265 by writing numbers in the empty boxes.



(2)

(d) The table below shows how the atomic radius of some atoms varies with atomic number.


Atomic number	Atomic radius in picometres (pm)
15	100
35	115
50	130
70	150
95	170

$$1 \text{ pm} = 10^{-12} \text{ m}$$

(i) On **Figure 2**, use the data from the table above to plot a graph of atomic radius against atomic number and draw a line of best fit.

Two points have been plotted for you.

**Figure 2**



(2)

- (ii) Scientists believe that the element with atomic number 126 can be produced and that it will be stable.

Use your graph in **Figure 2** to predict the atomic radius of an atom with atomic number 126.

Atomic radius = \_\_\_\_\_ pm

(1)

(Total 20 marks)